



ARIZONA-SONORA DESERT MUSEUM
PLANT CARE INFORMATION

Virgin's Bower or Barba de Viejo
Clematis drummondii

DESCRIPTION: Virgin's Bower is a perennial woody vine, found in a variety of habitats in the southern plains and southwestern U.S. and northern Mexico; from washes in the lower desert up into the oak forests. It climbs by tendrils and clasping leaves to nearly 20 feet. Leaves are deciduous. The flowers have 4 white, petal-like sepals surrounding a cluster of white stamens. The female flowers are followed by fluffy, whitish-rose colored seed plumes. Flowering can occur any time from spring until fall, from March to October.

RECOMMENDED USE: A background accent, shade plant, enhanced desert landscape or xeriscape, attract wildlife.

(over)



ARIZONA-SONORA DESERT MUSEUM
PLANT CARE INFORMATION

Virgin's Bower or Barba de Viejo
Clematis drummondii

DESCRIPTION: Virgin's Bower is a perennial woody vine, found in a variety of habitats in the southern plains and southwestern U.S. and northern Mexico; from washes in the lower desert up into the oak forests. It climbs by tendrils and clasping leaves to nearly 20 feet. Leaves are deciduous. The flowers have 4 white, petal-like sepals surrounding a cluster of white stamens. The female flowers are followed by fluffy, whitish-rose colored seed plumes. Flowering can occur any time from spring until fall, from March to October.

RECOMMENDED USE: A background accent, shade plant, enhanced desert landscape or xeriscape, attract wildlife.

(over)

CULTURE:



Hardiness: Roots are hardy to at least 10°F.



Sun tolerance: Light to moderate shade.



Watering and feeding: Low to moderate, depending on soil type.



Soil requirements: Best in sandy loam.



Pruning: As needed to control.

CULTURE:



Hardiness: Roots are hardy to at least 10°F.



Sun tolerance: Light to moderate shade.



Watering and feeding: Low to moderate, depending on soil type.



Soil requirements: Best in sandy loam.



Pruning: As needed to control.